

The Chandra Source Catalog Release 2.0

Ian N. Evans¹, Christopher Allen¹, Craig S. Anderson¹, Jamie A. Budynkiewicz¹, Douglas Burke¹, Judy C. Chen¹, Francesca Civano¹, Raffaele D'Abrusco¹, Stephen M. Doe², Janet D. Evans¹, Giuseppina Fabbiano^{1,} Daniel G. Gibbs II¹, Kenny J. Glotfelty¹, Dale E. Graessle¹, John D. Grier¹, Nicholas Lee¹, John C. Houck¹, Jennifer Lauer¹, Omar Laurino¹, Nicholas Lee¹, J. Rafael Martinez-Galarza¹, Michael L. McCollough¹, Jonathan C. McDowell¹, Joseph B. Miller¹, Joseph B. M Charles Paxson¹, David A. Plummer¹, Francis A. Primini¹, Arnold H. Rots¹, Meth A. Sundheim¹, Michael S. Tibbetts¹, David W. Van Stone¹, and Panagoula Zografou¹

¹Smithsonian Astrophysical Observatory ²Formerly Smithsonian Astrophysical Observatory ³Northrop Grumman Mission Systems ⁴MIT Kavli Institute for Astrophysics and Space Research

Highlights

- Includes almost all ACIS and HRC-I imaging observations from the period 1999–2014 (~560 deg² sky coverage).
- Source detection is performed on stacked (co-added) observations obtained using the same instrument, and with pointings colocated within a 60" diameter circle (because the *Chandra* PSF varies significantly with off-axis angle).
- The detection limit is ~ 5 net counts for on-axis sources with low-background exposures (typically < ~ 15 ks).

Comparison With Release 1.1

	Release 1.1	Release 2.0
Number of Sources	106,586	315,875 *
Number of Detections	158,071	374,349 *
Observation Dates	1999–2009	1999–2014
Source Detection Performed On	Individual Observations	Stacked Observations
Number of Observations	5,110	10,382
Number of Stacks	N/A	7,287
Largest Stack	N/A	81 Observations
Longest Effective Exposure	190 ks	5.8 Ms
Limiting Source Counts (on-axis)	~10	~5

- Detection properties are evaluated at both the stack and single observation level, and are combined to compute "best estimate" properties for each source on the sky.
- Key databased properties include positions and position errors, significance, processing flags/codes, extent, multi-band aperture photometry (total and net counts, count rates, photon and energy fluxes, model fluxes [multiple spectral models]), hardness ratios, multiple spectral model fits, inter- and intra-observation temporal variability.
- Numeric properties have associated independent lower and upper confidence limits; most properties are evaluated in 5 energy bands for ACIS, 1 for HRC-I.
- ~25 million science-ready FITS data products (~32 TB), including merged detection lists, detection region event lists, exposure maps, responses, spectra, light curves, aperture photometry PDFs, position error MCMC draws, per-Bayesian blocks properties, limiting sensitivity maps.

* The number of sources and detections included in the CSC 2.0 release may be reduced by <0.1% following final quality assurance processing.

Compact Source Detections In The Core Of Sgr A*



Position error ellipses computed from position confidence MCMC draws





— 1 arcmin

CSC 2.0 includes ~3900 compact sources detected in this image of the central region of the deepest Sgr A* stacked image (71 observations totaling ~2.2 Ms). The entire Sgr A* ensemble includes ~16,000 compact sources that were detected in 379 stacks (534 observations).



Spectral model fits and model energy fluxes determined using multiple source models



Source extent and local PSF models for every detection and energy band



0.00000.0001 0.0002 0.0003 Source Flux S (photons cm⁻² s⁻¹)

Hardness ratios



Intra- and inter-observation variability measures and light curves



For more information about CSC 2.0, see <u>http://cxc.cfa.harvard.edu/csc2/</u>



Elapsed Time (ks)

Acknowledgement. This work has been supported by NASA under contract NAS 8-03060 to the Smithsonian Astrophysical Observatory for operation of the Chandra X-ray Center.

0.02